



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

VI. Let ABC be \triangle right-angled at C . Produce AC to some point as D . Draw DF perpendicular to AB , produced, and meeting CB , produced.

Employing notation similar to that used in V., and proceeding somewhat in the same manner, we find that this method also yields a large number of proofs, in fact the same number that we found in V.

[To be Continued.]




Fig. 6.

ARITHMETIC.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

56. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

A, B, and C can walk at the rate of $a=3$, $b=4$, and $c=5$ miles, per hour. They start from Washington, at $m=1$, $n=2$, and $p=3$ o'clock, P. M., respectively. When B overtakes A, he is ordered (by A) back to C. When will B and C meet? Suppose B had ordered A back to C, when would A and C meet? In case all three continue walking ahead, at what time will they meet?

Solution by P. S. BERG, Larimore, North Dakota.

Since B gains 1 mile in 1 hour on A, to gain 3 miles will require 3 hours, or it will be 5 o'clock and 12 miles from starting point when B and A meet. C has traveled 10 miles. Since B and C travel 9 miles in 1 hour, they will travel 2 miles in $\frac{2}{3}$ hour, hence they will meet at $5\frac{2}{3}$ o'clock. Since A and C travel 8 miles in 1 hour, they will travel 2 miles in $\frac{1}{4}$ hour, hence they will meet at $5\frac{1}{4}$ o'clock.

In case all three continue walking ahead, as stated above A and B will meet at 5 o'clock. Since C gains 2 miles on A in 1 hour, to gain 6 miles will require 3 hours. Hence they will meet at 6 o'clock. Since C gains 1 mile on B in 1 hour, to gain 4 miles will require 4 hours. Hence it will be 7 o'clock when they meet.

Also solved by B. F. YANNEY and H. C. WILKS.

57. Proposed by L. B. FRAKER, Weston, Ohio.

Suppose that in a meadow the grass is of uniform quality and growth and that 6 oxen or 10 colts could eat up 3 acres of the pasture in $\frac{15}{16}$ of the time in which 10 oxen and 6 colts could eat up 8 acres; or that 600 sheep would require $2\frac{7}{8}$ weeks longer than 660 sheep to eat up 9 acres.